



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|---|--|---|--|
| (51) International Patent Classification 7 : A61K 47/48 | | A2 | (11) International Publication Number: WO 00/45849 |
| | | | (43) International Publication Date: 10 August 2000 (10.08.00) |
| (21) International Application Number: PCT/US00/02943 | | (81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). | |
| (22) International Filing Date: 4 February 2000 (04.02.00) | | | |
| (30) Priority Data: 60/118,802 5 February 1999 (05.02.99) US | | Published Without international search report and to be republished upon receipt of that report. | |
| (71) Applicants (for all designated States except US): GENZYME CORPORATION [US/US]; One Mountain Road, Framingham, MA 01701-9322 (US). STRESSGEN BIOTECHNOLOGIES CORPORATION [CA/CA]; 350-4243 Glanford Avenue, Victoria, British Columbia V8Z 4B9 (CA). | | | |
| (72) Inventors; and (75) Inventors/Applicants (for US only): SCHEULE, Ronald, K. [US/US]; 26 East Street, Hopkinton, MA 01748 (US). YEW, Nelson, S. [US/US]; 25 Rockdale Hill Circle, West Upton, MA 01568 (US). MIZZEN, Lee [CA/CA]; 1936 Quamichan Street, Victoria, British Columbia V8S 2C4 (CA). KADHIM, Salam, Abdul [CA/CA]; Unit 66, 14 Erskine Lane, Victoria, British Columbia V8Z 7J7 (CA). | | | |
| (74) Agents: GARRETT, Arthur, S.; Finnegan, Henderson, Farabow, Garrett & Dunner, L.L.P., 1300 I Street N.W., Washington, DC 20005-3315 (US) et al. | | | |

(54) Title: USE OF CATIONIC LIPIDS TO GENERATE ANTI-TUMOR IMMUNITY

(57) Abstract

A method of generating an anti-tumor immune response using a cationic molecule:biologically active molecule complex is provided. In one embodiment, the anti-tumor immune response is a protective, memory-based response. The complex may be administered alone, as the active ingredient in a formulation, or as an adjuvant. The invention also provides for methods of generating an immunostimulatory response against the tumor cell present during treatment by exposing a cationic molecule:biologically active molecule complex to a mammalian cell or a foreign tumor cell.

